



E445
JACC March 12, 2013
Volume 61, Issue 10



Congenital Cardiology Solutions

THE EFFECT OF MODIFIED ULTRAFILTRATION ON ANGIOPOIETINS IN PEDIATRIC CARDIOTHORACIC SURGERY

Poster Contributions

Poster Sessions, Expo North

Saturday, March 09, 2013, 10:00 a.m.-10:45 a.m.

Session Title: Congenital Cardiology Solutions: Congenital Surgical Therapies

Abstract Category: 13. Congenital Cardiology Solutions: Pediatric

Presentation Number: 1119-133

Authors: Sean Lang, Mansoor Syed, James Dziurja, Vineet Bhandari, John Giuliano, Jr., Yale School of Medicine, New Haven, CT, USA

Background: Cardiopulmonary bypass (CPB) subjects a patient's blood to hemodilution and nonphysiologic conditions which result in a systemic inflammatory response. Modified ultrafiltration (MUF) counteracts hemodilution, but may also improve outcomes by pro-inflammatory cytokine removal. Vascular growth factors angiotensin-1 (ang-1) and ang-2 play opposing roles in the capillary leak syndrome seen in patients following cardiac surgery. We hypothesize that the benefits of MUF include the removal of pro-inflammatory mediators, such as ang-2.

Methods: We performed a prospective cohort study. All patients ≤ 18 years of age undergoing cardiac surgery with CPB were offered enrollment. Three serum samples were obtained from each patient: 1. preoperatively, 2. following CPB, and 3. upon intensive care unit (ICU) admission. A final fluid sample from the MUF effluent was also analyzed. Ang-1 and ang-2 levels were determined using sandwich ELISA. The MUF effluent was used to calculate ang-1 and ang-2 percent extraction. The serum sample trends were analyzed by repeated measures ANOVA and paired t tests. The percent extraction was analyzed using the Wilcoxon Signed Rank Test.

Results: To date, 24 subjects have been enrolled. Mean ang-1 levels significantly decreased across all time points (pre CPB $2,987 \pm 3,012$ ng/mL, post CPB $1,635 \pm 1,046$ ng/mL, ICU admission $1,036 \pm 559$ ng/mL; $p < 0.01$). Ang-2 levels were significantly elevated at ICU admission when compared to both pre and post CPB levels (pre CPB $7,425 \pm 3,811$ ng/mL, post CPB $6,430 \pm 3,373$ ng/mL, ICU admission $11,126 \pm 7,914$ ng/mL; $p < 0.01$). There was no significant difference between the mean ang-1 or ang-2 percent extraction within MUF effluent ($1.8\% \pm 0.1\%$, $1.1\% \pm 3.2\%$, respectively; $p = 0.34$). In addition, the ang-2/1 ratio significantly increased across all time points (3.65 ± 2.41 , 5.84 ± 6.58 , 16.16 ± 17.87 , respectively; $p < 0.01$).

Conclusion: Ang-2/1 ratios significantly increase following CPB in children. The process of MUF removes both ang-1 and ang-2 equally. While data collection is ongoing, our preliminary results suggest that the clinical benefits of MUF cannot be attributed to the removal of larger quantities of ang-2 compared to ang-1.